

GLUEBALLS FROM IMPROVED LATTICE ACTIONS

COLIN MORNINGSTAR^a and MIKE PEARDON^b

^a *University of California at San Diego, La Jolla, CA 92093-0319*

^b *University of Kentucky, Lexington, KY 40506-0055*

The low-lying glueball masses and the hadronic scale r_0 are computed in lattice SU(3) gauge theory with the aim of establishing the effectiveness of the improved action approach in removing finite-spacing artifacts. The use of anisotropic lattices in which the temporal spacing is much smaller than that in the spatial directions allows much more efficient glueball mass measurements.

Glueballs and hybrid mesons are presently of great interest theoretically and experimentally. The lattice formulation of QCD provides an ideal setting in which to carry out theoretical studies of such systems from first principles using sophisticated numerical simulations. In order to extract the physical properties of glueballs and hybrid mesons from such simulations, systematic errors from the finite lattice spacing a must be removed or made acceptably small. There are two approaches to accomplishing this: (1) using finer grids or (2) using improved actions on coarse grids. The first approach is much simpler and has been used in almost all previous glueball and hybrid meson studies. However, this approach requires vast computational power. As the grid is made finer, many more lattice sites are needed to maintain the physical volume of the system. The simulation costs rise typically as $1/a^6$ as a is decreased. Because of this, lattice studies of glueballs have in the past been dominated by large collaborations using some of the world's fastest supercomputers.

Here, we show that the second approach, the use of improved actions, can be used to study glueballs much more efficiently. Improved actions have smaller lattice spacing errors, and hence, permit the use of much coarser lattices which can be simulated using modern computer workstations. The key to the success of the improvement program is the reliable determination of the couplings of the interactions terms in the action. Much effort over the past decade has been directed towards this problem. Recently, two competing methods have emerged, one which uses block-renormalization group transformations, another which advocates a judicious combination of mean field theory and perturbation theory. In this work, we use the latter approach.

A novel feature of our calculations is the use of anisotropic lattices in which the temporal spacing a_t is much smaller than the spacing a_s in the spatial directions. This allows much more efficient glueball mass measurements by exploiting the enhanced signal-to-noise of the glueball correlation functions at




Figure 1: Low-lying glueball masses m_g against lattice spacing a_s in terms of the hadronic scale $r_0 \approx 1/2$ fm defined by $[r^2 dV(r)/dr]_{r=r_0} = 1.65$ where $V(r)$ is the static quark potential. The $\times$ denote results for the scalar and tensor glueballs using the simple Wilson action¹. Results using the improved action for the A_1^{++} , E^{++} , T_2^{++} , and T_1^{+-} representations are indicated by $\circ$, $\square$, $\diamond$, and $*$, respectively.

smaller temporal separations. Mean-field link renormalization² is crucial for maintaining the proper renormalized anisotropy a_t/a_s .

Our results are shown in Fig. 1. The scalar glueball mass from the improved action exhibits dramatically reduced cutoff contamination compared to the Wilson action. Finite- a_s errors are seen to be small for the tensor and pseudo-vector glueballs, although differences between the E^{++} and T_2^{++} representations indicate small violations of rotational invariance, especially for large a_s . These results clearly show that glueballs can be studied without the use of supercomputers, provided that simulations are carried out using improved actions on anisotropic lattices.

References

1. J. Sexton, *et al.*, *Phys. Rev. Lett.* **75**, 4563 (1995); P. De Forcrand *et al.*, *Phys. Lett. B* **152**, 107 (1985); C. Michael and M. Teper, *Nucl. Phys. B* **314**, 347 (1989); UKQCD Collaboration, *Phys. Lett. B* **309**, 378 (1993).
2. G.P. Lepage and P.B. Mackenzie, *Phys. Rev. D* **48**, 2250 (1993).